



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AN ATTEMPT TO MODIFY THE AGGLUTINABILITY OF THE TYPHOID BACILLUS BY SELECTIVE ISOLATION OF INDIVIDUAL BACILLI.*

V. H. MOON.

(From the University of Kansas.)

All observers have noted in performing agglutination experiments, even with highly agglutinative serum in low dilutions, that some individual bacilli may resist agglutination and remain floating free in the serum. The object of this series of experiments was to determine whether this failure to clump is due to some peculiar property of the bacillus which can be transmitted to its descendants.

METHOD.

To isolate individual bacilli selectively seemed necessary as a means for this investigation. The most precise method for accomplishing this selective isolation is the technic designed and developed by Dr. M. A. Barber of the University of Kansas.¹ By using a microscopic capillary pipette supported in a mechanical holder provided with motion in three directions, one is able after considerable practice to isolate individual micro-organisms from a hanging drop. This drop is supported by a cover-glass over a moist chamber which is held and moved about under the microscope by the ordinary mechanical stage. The details of the method are fully described in the articles above referred to.

The medium used was meat broth carefully made according to the American Public Health Association standard, reaction 0.7 per cent acid to phenolphthalein. To begin with, an absolutely pure strain of *B. typhosus* was obtained by isolating an individual bacillus from a culture obtained from the laboratories of Johns Hopkins University. The resulting colony was transferred to broth and several agar slants were inoculated from this and labeled "Stock." A drop of the broth culture was placed on a sterile

* Received for publication April 1, 1911.

¹ *Jour. Infect. Dis.*, 1908, 5, p. 380; *Kansas University Science Bulletin*, 1907, p. 3.

cover-glass, and agglutination performed by adding an equal part of serum diluted 1:16. The serum used throughout this experiment was human serum from the same individual drawn under aseptic conditions and proved sterile by broth culture. It agglutinated typhoid bacilli at 1:96 dilution. As soon as clumping began a single definite clump was selected and isolated by means of the capillary pipette. The experiment was then allowed to stand one hour and 45 minutes at room temperature about 26° C. Then several individual bacilli which had resisted clumping were isolated and planted separately in small hanging drops. Chemically clean glass tubing was used for making all pipettes. For making each transfer of cultures, broth, or serum, and for each operation of isolation a new, freshly drawn pipette was used. The cover-glass was then transferred to a vaseline-sealed hollow-ground slide of large size and incubated 24 hours at 35° C. From this time on a double series of isolations and cultures was made. In the one series the first clumps to form were isolated, and in the other the individuals which resisted agglutination were selected and allowed to form colonies. For convenience we will refer to these as +S and -S respectively, the +S referring to that series selected on the basis of ready clumping and the -S to the series developed by selecting the resistant individuals. A number following the S indicates the number of selections made in that series.

After making five selections and cultures of the +S and four of the -S a preliminary comparative agglutination test was made with results as shown in the following table.

Dilution	1/16	1/24	1/32	1/48	1/64	1/96	Control
+S ₅	+	+	+	+	+	?	-
-S ₄	+	+	+	+	-	-	-

In making this test 24-hour agar cultures were used and the dilutions were examined microscopically. The writer in this case knew the identity of the cultures, but in all subsequent tests the cultures were marked in cipher by a second person and their identity revealed only after all observations had been made and recorded. One element of the personal equation was thus elimi-

Dilution	1/32	1/48	1/64	1/96	1/128	1/192	1/250	Control
+S ₉	+	+	+	+	+	+	—	—
—S ₈	+	+	?	—	—	—	—	—
Stock.....	+	+	+	+	—	—	—	—

Following this test the routine of performing agglutinations and selections was continued until 24 selections in the +S and 13 in the -S had been made. The average time of making the + selection was 20 minutes, while the - series were exposed to the action of the serum for two hours before the selections were made. Frequent transfers were made to clean freshly sterilized cover-glasses.

[illegible]

Following this test the cultures were kept at room temperature and transferred to fresh media three times in 18 days. Then a final test was made macroscopically, using both rabbit serum and the same human serum which was used in developing the two series. The cultures were marked in cipher as before.

[illegible]

HUMAN SERUM.

Dilution	1/16	1/24	1/32	1/48	1/64	1/96	1/128	Control
+S ₂₄	++	++	++	++	+	+	-	-
-S ₁₁	++	++	++	++	++	++	+	-
Stock.....	++	++	++	+	+	+	+	-

CONCLUSIONS.

The first two tests showed decided differences in the agglutinability of the two series. However in the final tests no such difference was evident. In the one the $-S_{13}$ agglutinated more readily than the $+S_{24}$, and in the other test the $-S_{13}$ agglutinated more readily than the stock. So we must conclude that, whatever differences in agglutinability were developed, they failed to become permanent or even to persist through the number of generations which would take place in an agar medium in 18 days at 26° C.

The author wishes to express his indebtedness to Dr. M. A. Barber for his patient instruction and oversight during the progress of the work.